



Elliott Bay Kidney Center - Center Hemodialysis
Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)
Through period ending Dec. 31, 2002

I. ADEQUACY OF DIALYSIS


	J	F	M	A	M	J	J	A	S	O	N	D
# eKt/V	95	92	90	94	91	92	88	93	92	91	94	95
# eKt/V >= 1.2	78	75	71	79	68	72	76	76	74	73	81	76
Pop. Mean =	1.44	1.36	1.41	1.40	1.33	1.39	1.39	1.42	1.38	1.39	1.43	1.38
Std. Deviation =	0.62	0.25	0.38	0.31	0.25	0.39	0.29	0.40	0.26	0.47	0.30	0.23
Network Mean =												
Natl. Mean =												
% eKt/V >= 1.2	82%	82%	79%	84%	75%	78%	86%	82%	80%	80%	86%	80%



# URR	95	92	90	94	91	93	89	94	93	95	96	95
# URR >= 65.0	81	83	79	86	79	81	81	88	85	84	85	88
Pop. Mean =	73.0	72.7	73.4	73.4	72.1	72.7	73.3	73.3	73.2	72.6	73.8	73.4
Std. Deviation =	12.3	8.9	7.4	5.3	6.5	6.8	6.2	6.1	6.1	8.0	6.3	5.62
*Network Mean =	72.4	72.4	72.4	72.4	72.4	72.4	72.4	72.4	72.4	72.4	72.4	72.4
Natl. Mean =	68.2	68.2	68.2	68.2	68.2	68.2	68.2	68.2	68.2	68.2	68.2	68.2
% URR >= 65.0	85%	90%	88%	91%	87%	87%	91%	94%	91%	88%	89%	93%
*Network % > 65.0	87%	87%	87%	87%	87%	87%	87%	87%	87%	87%	87%	87%
*Natl. % > 65.0	82%	82%	82%	82%	82%	82%	82%	82%	82%	82%	82%	82%



# Kt/V	95	92	90	94	91	92	88	93	92	91	94	95
# Kt/V >= 1.4	76	73	68	78	66	67	71	73	70	69	78	72
Pop. Mean =	1.66	1.56	1.62	1.61	1.53	1.60	1.61	1.63	1.58	1.60	1.64	1.59
Std. Deviation =	0.73	0.29	0.42	0.36	0.28	0.45	0.38	0.46	0.30	0.58	0.35	0.27
Network Mean =	1.61	1.61	1.61	1.61	1.61	1.61	1.61	1.61	1.61	1.61	1.61	1.61
Natl. Mean 1999 =	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40
% Kt/V >= 1.4	80%	79%	76%	83%	73%	73%	81%	78%	76%	76%	83%	76%
% Kt/V > 1.2	84%	93%	91%	95%	88%	92%	95%	97%	92%	89%	96%	95%
Network > 1.2 '00	93%	93%	93%	93%	93%	93%	93%	93%	93%	93%	93%	93%
Natl. % > 1.2 - 2001	91%	91%	91%	91%	91%	91%	91%	91%	91%	91%	91%	91%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.

Elliott Bay Kidney Center - Center Hemodialysis


Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)
Through period ending Dec. 31, 2002

I. ADEQUACY OF DIALYSIS - CONT.

J	F	M	A	M	J	J	A	S	O	N	D
<u>Blood Pressure Control (Pre)</u>											


# Pre Dia BP	96	93	93	94	93	93	91	94	94	95	96	95
# Pre Dia BP < 90	72	72	71	67	71	66	68	70	67	65	67	66
Pop. Mean =	82	80.5	81.7	83.3	81	81.2	80.3	80.9	81.4	81.7	81.8	82
Std. Deviation =	13.1	13.7	13.2	15.9	12.9	14.0	13.1	13.0	13.4	14.6	14.0	13.8
Network Mean =												
Natl. Mean =												
% Pre Dia BP < 90	75%	77%	76%	71%	76%	71%	75%	74%	71%	68%	70%	69%

% Pre Dia BP < 90




# Pre Sys BP	96	93	93	94	93	93	91	94	94	95	96	95
# Pre Sys BP < 150	51	59	49	52	53	57	47	52	50	52	51	50
Pop. Mean =	147	144	147	146	145	145	145	147	146	147	147	148
Std. Deviation =	21.6	20.9	22.1	19.9	19.2	20.0	20.0	20.8	19.0	20.4	21.3	18.5
Network Mean =												
Natl. Mean =												
%Pre Sys BP <150	53%	63%	53%	55%	57%	61%	52%	55%	53%	55%	53%	53%

% Pre Sys BP < 150



# Post/Dwt	92	94	93	93	91	94	94	95	95	95	95	95
# +/- 1% Dwt	59	52	50	57	54	58	65	63	64	57		
% +/- 1% of Dwt	64%	55%	54%	61%	59%	62%	69%	66%	67%	60%		

% +/- 1 % of Dwt



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)

Through period ending Dec. 31, 2002

I. ADEQUACY OF DIALYSIS - CONT.

	J	F	M	A	M	J	J	A	S	O	N	D
# CO2	96	92	93	94	93	93	91	94	94	95	96	95
# CO2 >= 23	46	41	43	49	28	51	58	36	53	52	46	50
Pop. Mean =	22	22	22	23	21	23	23	22	23	23	22	22
Std. Deviation =	2.60	3.27	2.46	2.39	2.45	2.49	2.16	2.32	2.68	2.48	2.62	3.07
Network Mean =												
Natl. Mean =												
% CO2 >= 23	48%	45%	46%	52%	30%	55%	64%	38%	56%	55%	48%	53%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis
Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)
Through period ending Dec. 31, 2002

*Albumin values use BCG method - as of July 2001


II. NUTRITION

	J	F	M	A	M	J	J	A	S	O	N	D
# Albumin	96	92	93	94	93	93	91	94	94	95	96	95
# Albumin >= 3.5	68	74	67	70	69	68	80	73	76	75	76	65
Pop. Mean =	3.52	3.67	3.6	3.63	3.6	3.63	3.7	3.68	3.72	3.68	3.71	3.6
Std. Deviation =	0.35	0.35	0.34	0.36	0.3	0.3	0.28	0.34	0.33	0.36	0.35	0.34
Network Mean =	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7
Natl. Mean =	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8
% Alb >= 3.5	71%	80%	72%	74%	74%	73%	88%	78%	81%	79%	79%	68%
EBKC % >= 3.7	36%	59%	51%	50%	46%	54%	52%	59%	62%	60%	63%	55%
EBKC % >= 4.0	8%	16%	15%	18%	12%	13%	21%	23%	26%	22%	23%	13%
*Natl % >= 3.5-BCG	80%	80%	80%	80%	80%	80%	80%	80%	80%	80%	80%	80%
*Natl. % >= 4.0-BCC	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%
*Netwk%>=3.5 BCC	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%
*Netwk%>=4.0 BCC	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%	29%

*Figures for 2000



# PCR/DPI	95	92	90	94	91	92	88	93	92	91	94	95
# PCR/DPI >= 1.2	27	29	29	25	29	24	33	25	25	26	30	27
Pop. Mean =	1.00	1.02	1.04	0.98	1.01	1.01	1.04	1.02	1.00	1.01	1.04	0.99
Std. Deviation =	0.29	0.29	0.28	0.26	0.26	0.28	0.24	0.27	0.27	0.27	0.29	0.27
Network Mean =												
Natl. Mean =												
% PCR/DPI >= 1.2	28%	32%	32%	27%	32%	26%	38%	27%	27%	29%	32%	28%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)


Through period ending Dec. 31, 2002

III. OSTEODYSTROPHY


	J	F	M	A	M	J	J	A	S	O	N	D
# PO4	96	92	93	94	93	93	91	94	94	95	96	95
# PO4 < 6.0	68	64	70	69	72							
# PO4 < 5.5						56	63	64	57	64	72	66
Pop. Mean =	5.3	5.2	5.0	5.1	5.1	5.2	4.9	5.0	5.2	4.9	4.7	4.7
Std. Deviation	1.49	1.53	1.6	1.44	1.53	1.57	1.33	1.44	1.54	1.44	1.58	1.44
Network Mean =												
Natl. Mean =												
% PO4 < 6.0 or 5.5	71%	70%	75%	73%	77%	60%	69%	68%	61%	67%	75%	69%



# PO4	96	92	93	94	93	93	91	94	94	95	96	95
# PO4 > 3.0	92	85	85	85	90							
# PO4 > 3.5						81	78	76	82	79	77	72
Pop. Mean =	5.3	5.2	5.0	5.1	5.1	5.2	4.9	5.0	5.2	4.9	4.7	4.7
Std. Deviation	1.49	1.53	1.6	1.44	1.53	1.57	1.33	1.44	1.54	1.44	1.58	1.44
Network Mean =												
Natl. Mean =												
% PO4 > 3.0 or 3.5	92%	88%	94%	94%	92%	87%	86%	81%	87%	83%	80%	76%



# Calcium	96	92	93	94	93	93	91	94	94	95	96	95
# Calcium < 11.0	93											
# Calcium < 10.2		79	84	77	82	80	80	82	78	85	78	85
Pop. Mean =	9.2	9.2	9.2	9.4	9.3	9.3	9.4	9.3	9.4	9.3	9.3	9.1
Std. Deviation	0.90	0.86	0.83	0.75	0.75	0.75	0.71	0.82	0.87	0.82	0.88	0.9
Network Mean =												
Natl. Mean =												
% Ca < 11.0 or 10.2	97%	86%	90%	82%	88%	86%	88%	87%	83%	89%	81%	89%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)


Through period ending Dec. 31, 2002

III. OSTEODYSTROPHY - CONT.


	J	F	M	A	M	J	J	A	S	O	N	D
# Calcium	96	92	93	94	93	93	91	94	94	95	96	95
# Calcium > 8.5	72	74	74	80	79	80	84	77	82	79	82	70
Pop. Mean =	9.2	9.2	9.2	9.4	9.3	9.3	9.4	9.3	9.4	9.3	9.3	9.1
Std. Deviation =	0.90	0.86	0.83	0.75	0.75	0.75	0.71	0.82	0.87	0.82	0.88	0.9
Network Mean =												
Natl. Mean =												
% Calcium > 8.5	75%	80%	80%	85%	85%	86%	92%	82%	87%	83%	85%	74%



# PTH-I	95	68	91	34
# PTH-I > 100	74	60	75	30
Pop. Mean =	314	393	457	422
Std. Deviation =	297	428	478	403
Network Mean =				
Natl. Mean =				
% PTH-I > 100	78%	88%	82%	88%



# PTH-I	95	68	91	34
# PTH-I < 300	56	37	45	17
Pop. Mean =	314	393	457	422
Std. Deviation =	297	428	478	403
Network Mean =				
Natl. Mean =				
% PTH-I < 300	59%	54%	49%	50%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)

Through period ending Dec. 31, 2002

III. OSTEODYSTROPHY - CONT.

	J	F	M	A	M	J	J	A	S	O	N	D
# Ca x PO4	96	92	93	94	93	93	91	94	94	95	96	95
# Ca x PO4 <= 65	82	79	82	85	82							
# Ca x PO4 <= 60						77	80	78	80	79	84	86
Pop. Mean =	48.7	48.1	45.2	47.4	47.8	47.8	45.7	46.1	48.6	45.1	44.1	43.2
Std. Deviation =	14.2	14.4	14.5	13.7	14.1	14.7	12.3	13.3	14.9	14.3	15	14.4
Network Mean =												
Natl. Mean =												
CaxPO4 <= 65or60	85%	86%	88%	90%	88%	83%	88%	83%	85%	83%	88%	91%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.


Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)


Through period ending Dec. 31, 2002

IV. ACCESS


	J	F	M	A	M	J	J	A	S	O	N	D
# Access	96	93	93	94	93	93	91	94	94	95	95	95
# Fistula	52	49	48	47	48	48	46	47	48	47	46	47
Network %												
Natl. % - 2000	30%	30%	30%	30%	30%	30%	30%	30%	30%	30%	30%	30%
% Fistula	54%	53%	52%	50%	52%	52%	51%	50%	51%	49%	48%	49%



# Access	96	93	93	94	93	93	91	94	94	95	95	95
# Graft	40	40	40	41	42	41	40	39	39	41	40	40
Network %												
Natl. %												
% Graft	42%	43%	43%	44%	45%	44%	44%	41%	41%	43%	42%	42%



# Access	96	93	93	94	93	93	91	94	94	95	95	95
# Other	4	4	5	6	3	4	5	8	7	7	9	8
Network %												
Natl. %												
% Other	4%	4%	5%	6%	3%	4%	5%	9%	7%	7%	9%	8%



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.

Elliott Bay Kidney Center - Center Hemodialysis

Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)

Through period ending Dec. 31, 2002


*Hct. values use Coulter method - as of July 2001

V. ANEMIA

	J	F	M	A	M	J	J	A	S	O	N	D
# Hct	96	93	93	94	91	92	91	94	93	95	96	94
# Hct > 30.0	92	87	84	85	82	84	83	87	85	91	89	85
Pop. Mean =	37.3	36.8	36.1	36.0	35.4	35.6	35.9	36.3	36.4	36.2	35.3	35.6
Std. Deviation =	4.53	4.74	4.6	4.56	4.51	4.21	4.25	4.57	4.82	4.32	4.04	4.28
Network Mean =	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9
Natl. Mean =	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2
% Hct > 30.0	96%	94%	90%	90%	90%	91%	91%	93%	91%	96%	93%	90%
*% Network > 30.0	89%	89%	89%	89%	89%	89%	89%	89%	89%	89%	89%	89%
% Natl. > 30.0	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%	79%

* Figures for 2000


% Hct > 30.0



# Hct	96	93	93	94	91	92	91	94	93	95	96	94
# Hct > 33.0	83	77	70	72	67	69	72	73	71	73	66	70
Pop. Mean =	37.3	36.8	36.1	36.0	35.4	35.6	35.9	36.3	36.4	36.2	35.3	35.6
Std. Deviation =	4.53	4.74	4.6	4.56	4.51	4.21	4.25	4.57	4.82	4.32	4.04	4.28
Network Mean =	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9	33.9
Natl. Mean =	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2	34.2
% Hct > 33.0	86%	83%	75%	77%	74%	75%	79%	78%	76%	77%	69%	74%
*% Network > 33.0	74%	74%	74%	74%	74%	74%	74%	74%	74%	74%	74%	74%
*% Natl. > 33.0	73%	73%	73%	73%	73%	73%	73%	73%	73%	73%	73%	73%

*Figures for 2000


% Hct > 33.0



# Hgb	92	91	94	93	95	96	95
# Hgb > 10.0	82	81	87	86	90	91	88
Pop. Mean =	11.8	11.8	12.1	12.1	12.1	11.9	12
Std. Deviation =	1.44	1.38	1.44	1.51	1.39	1.3	1.39
*Network Mean =	11.7	11.7	11.7	11.7	11.7	11.7	11.7
Natl. Mean =	11.7	11.7	11.7	11.7	11.7	11.7	11.7
% Hgb > 10.0	89%	89%	93%	92%	95%	95%	93%
*% Network >= 10	93%	93%	93%	93%	93%	93%	93%

*Figures for 2000

% Hgb > 10.0



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.

Elliott Bay Kidney Center - Center Hemodialysis


Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)

Through period ending Dec. 31, 2002

V. ANEMIA - CONT.

	J	F	M	A	M	J	J	A	S	O	N	D
# Hgb						92	91	94	93	95	96	95
# Hgb > 11.0						66	69	74	71	75	73	73
Pop. Mean =						11.8	11.8	12.1	12.1	12.1	11.9	12
Std. Deviation =						1.44	1.38	1.44	1.51	1.39	1.3	1.39
*Network Mean =	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7
Natl. Mean =												
% Hgb > 11.0						72%	76%	79%	76%	79%	76%	77%
*% Network >= 11	76%	76%	76%	76%	76%	76%	76%	76%	76%	76%	76%	76%


*Figures for 2000



Iron Profile


# Ferritin	94	93	90	95
# Ferritin > 100	80	81	79	86
Pop. Mean =	374	413	383	377
Std. Deviation =	347	368	463	308
*Network Mean =	489	489	489	489
Natl. Mean =				
% Ferritin > 100	85%	87%	88%	91%
*% Network > 100	88%	88%	88%	88%

*Figures for 2000



# Ferritin	94	93	90	95
# Ferritin < 800	86	86	85	90
Pop. Mean =	374	413	383	377
Std. Deviation =	347	368	463	308
*Network Mean =	489	489	489	489
Natl. Mean =				
% Ferritin < 800	91%	92%	94%	95%

*Figures for 2000



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.

Elliott Bay Kidney Center - Center Hemodialysis


Quality Assurance Reports (Excludes patients in first 3 months of treatment - as of Dec. 31, 2002)

Through period ending Dec. 31, 2002

V. ANEMIA - CONT.


	J	F	M	A	M	J	J	A	S	O	N	D
# % Sat	96			94			91			95		
# % Sat > 20	56			65			61			64		
Pop. Mean =	24.2			26.7			26.9			26.6		
Std. Deviation =	10.2			12.2			12.1			11		
*Network Mean =	28			28			28			28		
Natl. Mean =												
% Sat% > 20	58%			69%			67%			67%		
*% Network > 20	71%			71%			71%			71%		

*Figures for 2000



# % Sat	96	94	91	95
# % Sat < 50	92	91	85	92
Pop. Mean =	24.2	26.7	26.9	26.6
Std. Deviation =	10.2	12.2	12.1	11
*Network Mean =	28	28	28	28
Natl. Mean =				
% Sat% < 50	96%	97%	93%	97%

*Figures for 2000



*Methodology: a) Monthly values based on patient's monthly average of all values present for the month; b) Quarter values are average of all values for quarter; c) Values not averaged use last value present at months end.