Caring for Your IJ Catheter



Problem	What happens	What to do	How to prevent
Infection	 Fever, chills Redness, swelling, pus at catheter site Can spread to blood and heart; may cause death 	 Call your doctor right away Start antibiotics right away 	 Keep dressing and tape in place; staff will change dressing at dialysis Don't get catheter and dressing wet Touch the catheter as little as possible Get a fistula, graft, or PD catheter as soon as you can
Bleeding, catheter loose or comes out	Can be a medical emergency	 Press down with gauze or clean cloth until bleeding stops and call your doctor If it won't stop, or there's a lot of bleeding, call 911! Don't try to put it back in or move it Tape catheter down if you can 	 Don't allow clothing to rub or pull on catheter Keep dressing and tape in place Don't try to move catheter Get a fistula, graft, or PD catheter as soon as you can
Clotting Catheter clogs	May not be able to dialyze	May need to replace catheter	 Blood thinning medication (heparin) given with every treatment by staff Get a fistula, graft, or PD catheter as soon as you can



Caring for Your IJ Catheter, p. 2



Problem What happens What to do How to prevent	
Dressing gets wet, dirty or comes off **Risk of serious infection—blood and heart infection, and death* **Change dressing carefully (normally only staff should change dressing at dialysis) 1. Gather supplies: hand sanitizer, mask, tape, gauze 2x2 pad, Chlorascrub swab 2. Put on mask 3. Wash hands, then use hand sanitizer 4. Gently remove old tape and dressing 5. Use hand sanitizer again 6. Use Chlorascrub swab to clean, starting at catheter site and moving outward 7. Let skin dry 8. Cover with gauze 2x2 pad, holding pad by edges 9. Tape firmly	little as possible

